



The 2021
New Mexico
Alfalfa Variety
Test Report



Agricultural Experiment Station
**College of Agricultural, Consumer
and Environmental Sciences**

The 2021 New Mexico Alfalfa Variety Test Report

Leonard Lauriault, Ian Ray, Chris Pierce, Koffi Djaman, Robert Flynn, Mark Marsalis, Charles Havlik, Gasper Martinez, and Margaret West¹

Introduction

In 2021, 135,000 acres of alfalfa (*Medicago sativa*) were in production in New Mexico, which was a 20,000 acre decrease from 2020. Hay production was estimated at 689,000 tons. At a 2021 average of \$225 per ton (down from \$231 per ton in 2020), estimated gross returns from alfalfa hay produced in 2021 will total nearly \$155 million, which is a 2.6% increase from 2020. Besides its value for hay, alfalfa also is the legume of choice in irrigated perennial pastures. Whether used as pasture or hay, the value of alfalfa to New Mexico is greatly magnified by its contribution to livestock production and receipts from the sale of meat, milk, and other products generated by livestock enterprises.

Choosing a good alfalfa variety is a key step in establishing a highly productive stand of alfalfa, whether for hay or pasture. Differences between the highest- and lowest-yielding varieties in established irrigated tests included in this report ranged from 0.60 to 3.06 tons per acre in 2021. If sold as hay, this translates to a potential difference in returns of \$135 to \$689 per acre due to variety, or an increase of at least \$18 million for the industry in 2021 alone.

This report, which is a collaborative effort of New Mexico State University scientists at agricultural science centers throughout the state, provides yield data for alfalfa varieties included in yield trials in New Mexico. While consistently high yields compared to other varieties over a number of years and locations within a region is the best indication of varietal adaptation and persistence, other factors should be considered in the variety selection process (see NMSU's Cooperative Extension Service Circular 654, Selecting alfalfa varieties for New Mexico). In addition to fall dormancy and winter hardiness, high levels of pest resistance are critical to protecting an alfalfa stand for long-term production. Alfalfa grown in New Mexico should have at least a resistant (R) rating for bacterial wilt, Fusarium wilt, anthracnose, Phytophthora root rot, spotted alfalfa aphid, blue alfalfa aphid, pea aphid, stem nematode, and southern rootknot nematode. Seed quality also should be high. Selecting an alfalfa variety based on seed cost is a gamble producers often lose. To be assured of achieving a long-lasting, highly productive stand, buy either certified or Plant Variety Protected (PVP) seed, which guarantees the genetics and performance. The best choice of seed of any variety is one that was treated with a fungicide and nitrogen-fixing bacteria before it was bagged.

Description of Tests

Replicated alfalfa variety tests included in this report were conducted under research controls at NMSU's Agricultural Science Centers at Las Cruces (2019, full and summer terminated irrigation), Los Lunas (2019), Tucumcari (2018), and Farmington (2018 and 2020). Weather data for 2021 and the long-term averages from all locations are presented in Table 1.

Yield data (on a dry matter basis) are presented in Tables 2-7. Varieties are listed in order from highest to lowest average annual production. Yields are given by cutting for 2021 and by year for each production year. Statistical analyses were performed on all alfalfa yield data (including experimental entries) to determine if the apparent differences are truly due to variety or just to chance. The variety with the highest numerical yield in each column is marked with two asterisks (**), and those varieties not significantly different from that variety are marked with one asterisk (*). Those are the varieties from which to make an initial selection. Otherwise, to determine if two varieties are truly different, compare the difference between the two varieties to the Least Significant Difference (LSD) at the bottom of the column. If the difference is equal to or greater than the LSD, the varieties are truly different in yield when grown under the conditions at a given location. If NS is given for the LSD, there was no statistical difference between the highest and lowest yielding varieties. The Coefficient of Variation (CV), which is a measure of the variability of the data, is included for each column of means. Low variability (<20 percent) is desirable, and increased variability within a study results in higher CVs and larger LSDs. There might be a difference between previously published data and the data given in this publication for the same tests because of differences in the programs used for statistical analysis.

Table 8 summarizes information about proprietors, Roundup Ready genetics, fall dormancy, winter survival (measured in the northern United States), pest resistance, and yield performance across years and locations for all varieties currently included in NMSU's alfalfa variety testing program. For information about other varietal characteristics, such as grazing, salt, or traffic tolerance or GMO traits besides Roundup Ready® genetics, check the National Alfalfa and Forage Alliance website for the Alfalfa Variety Leaflet (<https://www.alfalfa.org/varietyLeaflet.php>). In Table 8, varieties are listed alphabetically by fall dormancy category. As in the data tables, the variety with the highest numerical yield in each column is marked with two asterisks (**), and those varieties not significantly different from that variety are marked with one asterisk (*). Remember good performance across several years and locations is the best indicator of broad adaptation, pest resistance, and persistence.

Seed labeled "common," "variety not stated," or "variety unknown", particularly that from other states, is of unknown genetic background and may or may not have the necessary disease or insect resistance. New Mexico Common and African Common seed used in all tests throughout the state has come from the same supplier and seed fields in New Mexico. Seed purchased from other dealers may or may not be of the same quality and performance.

Summary

Consistent production of high alfalfa yields is the result of selecting good varieties and implementing good management techniques. Soil fer-

¹Forage Crop Management Scientist and Superintendent, NMSU Rex E. Kirksey Agricultural Science Center at Tucumcari; Alfalfa Breeder, NMSU, Las Cruces; Forage Research Scientist, NMSU, Las Cruces; Agronomist, NMSU Agricultural Science Center at Farmington; Extension Agronomist and Superintendent, NMSU Agricultural Science Center at Artesia; Extension Forage Specialist and Superintendent, NMSU Agricultural Science Center at Los Lunas; Senior Research Assistant, NMSU Agricultural Science Center at Los Lunas; Research Assistant, NMSU Rex E. Kirksey Agricultural Science Center at Tucumcari; and Agricultural Research Scientist, NMSU Agricultural Science Center at Farmington; respectively.

tility should be maintained at recommended levels based on soil tests, irrigation should be properly applied, weeds and insects should be controlled using appropriate cultural and/or chemical methods, and harvest management should allow sufficient time to restock root energy prior to winter. For dormant (FD 1 to 3) and semidormant (FD 4 to 6) varieties, a 6-week rest period before a dormancy-inducing freeze (27°F) is recommended to allow plants to replenish root reserves for winter survival and initiate spring growth, after which harvesting might be done either mechanically or by grazing. Non-dormant (FD 7 to 9) varieties also might benefit from this rest period. Removing fall growth is beneficial to reducing weevil populations the following year as eggs are laid in and overwinter in stems. Harvesting established stands at early bloom would result in 3 to 5 cuttings per year before initiation of the rest period in most areas of New Mexico. More dormant varieties might not produce yields that can be baled during the rest period; however, these can still be grazed. For additional information about alfalfa management, refer to the other NMSU Agricultural Experiment Station and Cooperative Extension Service publications listed in Table 9.

Acknowledgements

The authors express appreciation to the following for their significant contribution to The New Mexico Alfalfa Variety Testing Program by helping with planting, maintaining, harvesting, or other data collection, and data entry: Dallen Begay, Jason Box, Patty Cooksey, Josh Foster, Lorenzo Gallegos, Ryan Garcia, Christopher Hill, Jared Jennings, Shane Jennings, Jonah Joe, Martin Lopez, Ruben Pacheco, Tom Place, Dennis Price, Patricia Cooksey, Franklin Jason Thomas, and the staff at University Marketing and Communications who make publications such as this possible. Salaries and research support were provided by state and federal funds appropriated to the New Mexico Agricultural Experiment Station. Federal funds included those appropriated through the Hatch Act of 1887.

Table 1. Temperature and precipitation data for 2021 and the long-term averages for the New Mexico Alfalfa Variety Test locations.

Location Elevation Latitude	Las Cruces ¹ 3832 ft. 32° 12' N				Los Lunas 4840 ft. 34° 46' N				Tucumcari 4091 ft. 35° 12' N				Farmington 5640 ft. 36° 41' N			
	Temp. (°F)		Precip. (in.)		Temp. (°F)		Precip. (in.)		Temp. (°F)		Precip. (in.)		Temp. (°F)		Precip. (in.)	
Month	2021	Avg.	2021	Avg.	2021	Avg.	2021	Avg.	2021	Avg.	2021	Avg.	2021	Avg.	2021	Avg.
Nov-20	52	49	0.02	0.53	48	44	0.06	0.46	45	47	0.54	0.68	44	41	0.16	0.65
Dec-20	39	41	0.22	0.68	34	35	0.25	0.52	42	39	0.12	0.64	27	31	0.19	0.47
Jan-21	40	40	0.12	0.56	35	35	0.22	0.38	39	38	0.18	0.40	30	30	0.50	0.51
Feb-21	47	45	0.07	0.37	39	40	0.46	0.42	36	42	0.10	0.50	35	36	0.44	0.48
Mar-21	54	52	0.00	0.22	46	47	0.12	0.46	50	49	1.52	0.76	41	43	0.93	0.64
Apr-21	64	60	0.09	0.21	56	55	0.32	0.48	57	57	0.13	1.11	52	51	0.08	0.58
May-21	71	69	0.00	0.29	65	64	0.00	0.45	67	66	2.12	1.86	61	60	0.07	0.53
Jun-21	79	78	1.40	0.72	77	73	0.78	0.56	78	76	1.22	1.85	73	70	0.29	0.26
Jul-21	78	80	2.99	1.36	77	77	2.44	1.38	77	79	6.51	2.63	77	76	0.35	0.81
Aug-21	77	78	2.23	2.29	75	75	1.21	1.62	79	77	2.19	2.67	73	74	0.68	1.05
Sep-21	71	71	0.48	1.38	70	68	0.44	1.15	75	71	0.55	1.54	66	66	0.77	1.10
Oct-21	60	61	0.02	0.91	54	56	0.39	1.04	62	59	0.98	1.33	52	54	0.85	0.97
Annual	61	60	7.64	9.52	56	56	6.69	8.92	59	58	16.16	15.97	53	53	5.31	8.05

¹Long-term averages for the Las Cruces test site are from the State University weather station, located approximately 5.5 miles to the north.

Table 2. Dry matter yields (tons/acre) of alfalfa varieties sown September 27, 2019, at NMSU's Leyendecker Plant Science Research Center at Las Cruces and flood-irrigated twice per cutting (normal irrigation)†.

Variety Name	2020	2021 Harvests							2021	2-Year
	Total	1-May	27-May	24-Jun	22-Jul	27-Aug	29-Sep	12-Nov	Total	Average
SW9812	6.05**	1.18*	1.08*	0.94*	0.91*	1.19*	0.95*	0.62*	6.83*	6.56**
SW9813	5.86*	1.14*	1.04*	0.97**	0.90*	1.17*	0.93*	0.66*	6.79*	6.37*
NM 1705PAR	5.56*	1.16*	1.09**	0.97**	1.01**	1.27**	0.97**	0.65*	7.11**	6.36*
NM 1701PAR	5.81*	1.17*	1.06*	0.92*	0.84	1.10	0.90*	0.57	6.52*	6.29*
NM 1703PAR	5.95*	1.13*	1.07*	0.94*	0.87	1.19*	0.92*	0.63*	6.76*	6.26
SW8888	5.73*	1.09*	1.01*	0.89*	0.88	1.17*	0.96*	0.63*	6.61*	6.24*
NM 1704PAR	6.01*	1.13*	1.02*	0.87*	0.85	1.12	0.87	0.54	6.40*	6.23*
Mesquite	5.78*	1.05*	1.01*	0.90*	0.85	1.16*	0.92*	0.63*	6.51*	6.23*
TMA 990	5.81*	1.05*	0.95*	0.90*	0.87	1.16*	0.90*	0.69**	6.52*	6.19*
AmeriStand 803T	5.91*	1.06*	1.02*	0.92*	0.85	1.10	0.89*	0.60*	6.45*	6.15*
SW1011	5.64*	1.12*	1.06*	0.96*	0.85	1.19*	0.92*	0.66*	6.78*	6.15*
SW9720	5.74*	1.15*	1.04*	0.94*	0.86	1.12	0.87	0.64*	6.65*	6.12*
Dona Ana	5.26*	1.20*	1.08*	0.96*	0.91*	1.20*	0.91*	0.62*	6.91*	5.97*
SW8421S	5.47*	1.01*	0.97*	0.90*	0.81	1.15	0.92*	0.61*	6.38*	5.91
NM 170506PAR	5.53*	1.07*	0.99*	0.89*	0.83	1.12	0.89*	0.60*	6.40*	5.90
AFX 779	5.58*	1.07*	0.99*	0.85*	0.77	1.07	0.87	0.56	6.18	5.84
NuMex Bill Melton	5.19*	1.21**	1.05*	0.90*	0.80	1.06	0.82	0.57	6.41*	5.82
NM 1702PAR	5.43*	0.99*	0.91*	0.78	0.72	0.99	0.79	0.52	5.70	5.63
AFX 149092	5.11*	0.99*	0.96*	0.87*	0.82	1.12	0.84	0.61*	6.25	5.61
NM 1712PAR	5.60*	1.02*	0.91*	0.77	0.73	0.95	0.74	0.40	5.50	5.54
NM 1713PAR	4.93*	0.95*	0.81*	0.69	0.67	0.85	0.71	0.34	5.01	5.02
NM 1715PAR	4.89*	0.97*	0.86*	0.69	0.67	0.89	0.68	0.33	5.10	4.88
Mean	5.58	1.09	1.00	0.88	0.83	1.11	0.87	0.58	6.35	5.97
LSD (0.05)	NS	NS	NS	0.14	0.13	0.12	0.10	0.11	0.85	0.63
CV%	9.80	12.15	11.43	11.13	10.96	7.80	8.27	13.59	9.51	10.56

†Data were analyzed with analysis of covariance using check plots of NuMex Bill Melton as the covariate. 2020 Harvest dates: 19-Jun, 22-Jul, 21-Aug, 15-Sep, 19-Oct, and 17-Nov.

**Highest numerical value in the column.

*Not significantly different from the second highest numerical value in the column based on the 5% LSD. NS means that there were no significant differences between the varieties within that column at the 5% level.

Table 3. Dry matter yields (tons/acre) of alfalfa varieties sown September 28, 2019, at NMSU's Leyendecker Plant Science Research Center at Las Cruces and flood-irrigated twice per cutting through June and resuming in mid-September through October (summer irrigation termination)†.

Variety Name	2020 Total	2021 Harvests						2021 Total	2-Year Average
		30-Apr	27-May	25-Jun	22-Jul	26-Aug	11-Nov		
SW8421S	4.03*	1.22	1.14	1.18**	0.92**	0.25*	0.63**	5.33*	4.68**
AFX 779	4.04*	1.32*	1.19*	1.17*	0.81	0.19	0.55	5.23*	4.64*
NM 1703PAR	3.80*	1.29*	1.15*	1.16*	0.91*	0.30**	0.57	5.39*	4.62*
NM 1704PAR	3.97*	1.35**	1.19*	1.15*	0.85*	0.20	0.55	5.29*	4.61*
Dona Ana	4.07**	1.20	1.12	1.09	0.79	0.24	0.63**	5.08*	4.60*
SW9813	3.89*	1.25*	1.14	1.16*	0.88*	0.25*	0.61*	5.28*	4.58*
SW9812	3.71*	1.33*	1.18*	1.16*	0.87*	0.24	0.63**	5.42**	4.57*
SW9720	3.86*	1.19	1.11	1.13*	0.87*	0.24	0.60*	5.14*	4.53*
TMA 990	3.60*	1.11	1.09	1.13*	0.88*	0.30**	0.79***	5.31*	4.47*
NM 170506PAR	3.60*	1.22	1.18*	1.16*	0.90*	0.26*	0.59*	5.31*	4.45*
AmeriStand 803T	3.69*	1.18	1.12	1.13*	0.85*	0.25*	0.59*	5.12*	4.42*
NM 1702PAR	3.62*	1.29*	1.19*	1.15*	0.82*	0.19	0.55	5.19*	4.41*
NM 1701PAR	3.47*	1.31*	1.22**	1.16*	0.89*	0.27*	0.55	5.41*	4.41*
NM 1705PAR	3.58*	1.26*	1.17*	1.14*	0.88*	0.22	0.54	5.22*	4.38*
SW8888	3.85*	1.17	1.08	1.06	0.83*	0.19	0.51	4.84	4.37*
SW1011	3.49*	1.18	1.10	1.13*	0.88*	0.26*	0.59*	5.14*	4.36*
AFX 149092	3.69*	1.14	1.11	1.11*	0.84*	0.24	0.60*	5.05*	4.34*
NuMex Bill Melton	3.66*	1.24	1.12	1.07	0.78	0.20	0.53	4.94	4.29
Mesquite	3.45*	1.18*	1.14	1.12*	0.84*	0.24	0.57	5.11*	4.25
NM 1712PAR	3.76*	1.30*	1.11	1.03	0.73	0.16	0.35	4.68	4.24
NM 1715PAR	3.27*	1.23	1.10	1.01	0.71	0.16	0.26	4.47	3.88
NM 1713PAR	3.42*	1.20	1.03	0.92	0.69	0.15	0.30	4.30	3.86
Mean	3.71	1.23	1.14	1.11	0.84	0.23	0.55	5.10	4.41
LSD (0.05)	NS	0.12	0.08	0.09	0.11	0.06	0.06	0.42	0.37
CV%	8.71	7.03	5.05	5.83	9.18	20.39	7.99	5.83	8.37

†Data were analyzed with analysis of covariance using check plots of NuMex Bill Melton as the covariate. 2020 Harvest dates, 18-Jun, 21-Jul, and 17-Oct.

***Highest numerical value in the column and significantly greater than all other values.

**Second highest numerical value in the column.

*Not significantly different from the second highest numerical value in the column based on the 5% LSD.

NS means that there were no significant differences between the varieties within that column at the 5% level.

Table 4. Dry matter yields (tons/acre) of flood-irrigated alfalfa varieties sown September 27, 2019, at NMSU's Agricultural Science Center at Los Lunas†.

Variety Name	2020 Total	2021 Harvests				2021 Total	2-Year Average
		26-May	12-Jul	26-Aug	21-Oct		
NM 1704PAR	11.97**	2.82*	2.58	1.89	1.33	8.61	10.29**
NM 1713PAR	11.42*	3.01*	2.63*	2.01*	1.42	9.06*	10.24*
NM 1712PAR	11.47*	3.01*	2.54	2.05*	1.31	8.91*	10.19*
AFX 779	11.71*	2.56	2.56	2.00*	1.50*	8.61	10.16*
NuMex Bill Melton	11.12*	2.73	2.67*	2.13*	1.46*	8.98*	10.05*
SW7473	11.06*	2.91*	2.68*	1.93	1.39	8.90*	9.98*
AmeriStand 715NTRR	11.32*	2.67	2.32	1.96*	1.57**	8.51	9.92*
Roadrunner	10.56	3.03**	2.84**	2.06*	1.35	9.27**	9.92*
NM 1715PAR	10.88	2.72	2.43	2.07*	1.46*	8.67	9.78*
Artesian Sunrise	10.53	2.67	2.72*	2.05*	1.34	8.77	9.65
Zia	10.91	2.61	2.42	1.82	1.35	8.20	9.56
WL 454HQ.RR	10.49	3.00*	2.41	1.78	1.38	8.57	9.53
NM Common	10.48	2.58	2.26	1.95	1.43	8.22	9.35
Artesian Sun 6.3	9.91	2.67	2.46	2.14**	1.28	8.55	9.23
WL 440HQ	10.16	2.69	2.40	1.91	1.29	8.28	9.22
Dona Ana	10.08	2.52	2.47	2.00*	1.33	8.31	9.19
AFX 579	10.06	2.69	2.25	1.93	1.30	8.17	9.11
SW3407	10.03	2.73	2.20	1.82	1.40	8.14	9.09
AFX 469	9.85	2.84*	2.26	1.88	1.19	8.16	9.01
Hi-Gest 660	9.76	2.56	2.23	1.91	1.30	8.00	8.88
SW4107	9.06	2.83*	2.56	1.79	1.31	8.49	8.77
SW5511	9.16	2.54	2.47	1.70	1.31	8.01	8.58
Mean	10.54	2.74	2.47	1.94	1.36	8.52	9.53
LSD (0.05)	1.06	0.24	0.26	0.20	0.13	0.43	0.61
CV%	7.09	6.32	7.42	7.11	6.65	3.56	6.40

†Data were detrended using nearest neighbor analysis and analyzed using analysis of variance.

2020 Harvest dates: 9-Jun, 16-Jul, 8-Sep, and 2-Nov.

**Highest numerical value in the column.

*Not significantly different from the second highest numerical value in the column based on the 5% LSD.

NS means that there were no significant differences between the varieties at the 5% level.

Table 5. Dry matter yields (tons/acre) of alfalfa varieties sown September 14, 2018, at NMSU's Rex E. Kirksey Agricultural Science Center at Tucumcari and not irrigated in 2021†.

Variety Name	2020 Total	2021 Harvests		2021 Total	2-Year Average
		28-Jul	21-Sep		
African Common	2.15*	1.33*	1.00**	2.34**	2.24**
SW7408	2.21**	1.41*	0.85*	2.26*	2.23*
Zia	2.11*	1.38*	0.86*	2.24*	2.18*
Dona Ana	2.00*	1.46**	0.87*	2.33*	2.16*
SW8476	2.06*	1.46**	0.67	2.13*	2.09*
SW8409	1.91*	1.38*	0.61	1.99	1.95
SW8412	1.96*	1.27*	0.62	1.90	1.93
NM Common	1.74	1.21*	0.79*	1.99	1.87
SW7473	1.79	1.23*	0.69	1.92	1.86
SW5210	1.86*	1.17*	0.62	1.79	1.82
HybriForce-3600	1.66	1.22*	0.75	1.97	1.82
Hi-Gest 660	1.69	1.29*	0.61	1.91	1.80
SW8421S	1.78	1.21*	0.59	1.80	1.79
SW1517	1.64	1.27*	0.60	1.87	1.75
SW1509	1.68	1.18*	0.53	1.71	1.70
Mean	1.88	1.30	0.71	2.01	1.95
LSD (0.05)	0.38	NS	0.23	0.34	0.28
CV%	14.01	13.90	22.97	11.99	14.09

†Data were detrended using nearest neighbor analysis and analyzed using analysis of variance.

No harvests were taken in 2019 due to harvesting equipment and irrigation system failures; Irrigation system failures also occurred in 2020 that were exacerbated by parts availability issues due to COVID-19. 2020 Harvest dates: 13-Jul and 21-Oct.

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

NS means that there were no significant differences between the varieties within that column at the 5% level.

Table 6. Dry matter yields (tons/acre) of sprinkler-irrigated alfalfa varieties sown August 28, 2018, at NMSU's Agricultural Science Center at Farmington†.

Variety Name	2019 Total	2020 Total	2021 Harvests			2021 Total	3-year Average
			17-Jun	8-Aug	21-Sep		
FSG 423ST	9.44**	11.52*	1.19*	5.48**	2.76*	9.43**	10.13**
SW3407	9.43*	11.93*	1.15*	4.51*	2.83*	8.48*	9.94*
6422Q	9.40*	11.88*	1.19*	4.38*	2.71*	8.27*	9.85*
SW5207	8.82*	12.63**	1.06*	4.09*	2.82*	7.97*	9.81*
WL 365HQ	9.09*	11.97*	1.35*	4.19*	2.64*	8.18*	9.75*
FSG 415BR	9.17*	12.17*	1.30*	3.82*	2.65*	7.76*	9.70*
SW4107	8.82*	11.69*	1.63**	4.18*	2.58*	8.39*	9.63*
Lahontan	8.48*	11.78*	1.03*	4.72*	2.74*	8.49*	9.59*
FSG 426	9.21*	11.09*	1.00*	4.42*	2.29*	7.71*	9.33*
FSG 524	8.22*	11.92*	1.27*	4.17*	2.36*	7.79*	9.31*
Ranger	9.51*	10.21*	1.17*	4.05*	2.63*	7.85*	9.19*
FSG 403LR	8.52*	10.94*	1.26*	3.71*	2.87**	7.84*	9.10*
6585Q	8.17*	11.57*	1.11*	3.77*	2.45*	7.33*	9.02*
WL 377HQ	8.37*	10.51*	1.22*	4.16*	2.48*	7.85*	8.91*
AmeriStand 518NT	8.04*	11.49*	1.04*	3.48*	2.49*	7.00*	8.84*
NM Common	8.19*	11.01*	1.03*	3.00*	2.34*	6.37*	8.52*
Mean	8.80	11.52	1.19	4.13	2.60	7.92	9.41
LSD (0.05)	NS	NS	NS	NS	NS	NS	NS
CV%	10.26	10.25	25.44	23.52	12.65	15.39	14.99

†Data were detrended using nearest neighbor analysis, and analyzed using analysis of variance.

2019 Harvest dates: 17-Jun, 24-Jul, and 3-Sep.

2020 Harvest dates: 9-Jun, 22-Jul, 26-Aug, and 7-Oct.

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

NS means that there were no significant differences between the varieties within that column at the 5% level.

Table 7. Dry matter yields (tons/acre) of sprinkler-irrigated alfalfa varieties sown August 10, 2020, at NMSU's Agricultural Science Center at Farmington†.

Variety Name	2021 Harvests			2021 Total
	19-Jun	10-Aug	22-Sep	
NM 1715PAR	0.60*	4.04**	2.75*	7.39**
Bar MSI	0.43*	3.65*	2.84**	6.92*
SW4412Y	0.76**	3.47*	2.62*	6.85*
SW3407	0.63*	3.34	2.84**	6.81*
Magnum 8	0.74*	3.44	2.61*	6.78*
African Common	0.74*	3.51*	2.44	6.68*
NM 1713PAR	0.49*	3.53*	2.66*	6.68*
NM 1712PAR	0.44*	3.31	2.58*	6.33
NM Common	0.66*	3.11	2.52*	6.29
SW4506	0.53*	3.15	2.54*	6.22
SW4107	0.54*	2.91	2.70*	6.14
Dona Ana	0.45*	3.03	2.64*	6.12
AFX 469	0.60*	3.12	2.35	6.06
Zia	0.44*	3.17	2.02	5.62
Mean	0.57	3.34	2.58	6.49
LSD (0.05)	NS	0.59	0.37	0.92
CV%	31.41	12.29	9.95	9.88

†Data were analyzed using analysis of variance.

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

NS means that there were no significant differences between the varieties within that column at the 5% level.

Table 9. New Mexico State University Cooperative Extension Service and Agricultural Experiment Station publications related to alfalfa management.

Number	Title
A-123	Sampling for plant tissue analysis
A-129	Nitrogen fixation by legumes
A-130	Inoculation of legumes
A-131	Certified seed
A-137	Soil analysis: A key to soil nutrient management
A-145	Certified noxious weed free program
A-146	Appropriate analysis for New Mexico soils
A-229	Phymatotrichum root rot
A-325	Managing weeds in alfalfa
A-326	Downy mildew on alfalfa
A-333	User manual of the alfalfa yield predictor
A-334	Beet armyworm in New Mexico Hay
A-335	Variegated cutworm in New Mexico Hay
A-336	Managing Roundup Ready alfalfa and conventional or organic alfalfa hay in nearby fields in New Mexico
A-337	Recommendations for Roundup Ready alfalfa weed management and stand removal in New Mexico
A-338	Alfalfa weevil control options in New Mexico
A-339	Alfalfa integrated pest management: Aphids
A-615	Dodder (<i>Cuscuta ssp.</i>) biology and management
H-158	How to collect and send plant specimens for disease diagnosis
CR-536	Blister beetles in alfalfa
CR-586	Irrigated pasture management in New Mexico
CR-633	Using a computer application to predict irrigated alfalfa yield
CR-641	Hay quality, sampling and testing
CR-644	Assessing alfalfa stands after winter injury, freeze damage, or any time renovation is considered in New Mexico
CR-646	Managing alfalfa during drought
CR-654	Selecting alfalfa varieties for New Mexico
CR-659	Whitefringed beetle in New Mexico alfalfa
CR-668	Reducing harvest and post-harvest losses of alfalfa and other hay
CR 682	GMO Crops in New Mexico agriculture
CR-696	Perennial forage species for irrigated pastures and hay in New Mexico

These publications, and alfalfa variety test reports from previous years, are available from your county office of the NMSU Cooperative Extension Service or online at <http://forages.nmsu.edu/resources.html> and aces.nmsu.edu/pubs/



New Mexico State University
BE BOLD. Shape the Future.

The College of Agricultural, Consumer and Environmental Sciences is an engine for economic and community development in New Mexico, improving the lives of New Mexicans through academic, research, and Extension programs.

New Mexico State University is an equal opportunity/affirmative action employer and educator. NMSU and the U.S. Department of Agriculture cooperating.